

## **ZOVIRAX<sup>®</sup> Cream for Recurrent Cold Sores**

ZOVIRAX<sup>®</sup> (acyclovir) Cream 5% is indicated for the treatment of recurrent herpes labialis (cold sores) in adults and adolescents (12 years of age and older).

### **Important Safety Information**

ZOVIRAX<sup>®</sup> Cream is intended for cutaneous use only and should not be used in the eye or inside the mouth or nose. ZOVIRAX<sup>®</sup> Cream has a potential for irritation and contact sensitization. In clinical trials, the most common adverse reactions at the site of topical application occurred in less than 1% of patients and included dry lips, desquamation, dryness of skin, cracked lips, burning skin, pruritus, flakiness of skin, and stinging on skin. ZOVIRAX<sup>®</sup> Cream does not prevent transmission of HSV infections, and its effect has not been established in immunocompromised patients.

**ZOVIRAX<sup>®</sup> Cream is available by prescription only.**

## **ZOVIRAX<sup>®</sup> Ointment for Initial Genital Herpes Outbreak**

ZOVIRAX<sup>®</sup> (acyclovir) Ointment 5% is indicated in the management of initial genital herpes and in limited non-life-threatening mucocutaneous herpes simplex virus infections in immunocompromised patients.

### **Important Safety Information**

ZOVIRAX<sup>®</sup> Ointment is intended for cutaneous use only and should not be used in the eye. Mild pain (including transient burning and stinging) has been reported at an incidence comparable to placebo (30%). ZOVIRAX<sup>®</sup> Ointment does not prevent the transmission of HSV infections and should not be used for the prevention of recurrent HSV infections.

**ZOVIRAX<sup>®</sup> Ointment is available by prescription only.**